

The very high redshift component of the OTELO survey

Ángel Bongiovanni^{1,2}, Marina Ramón-Pérez^{1,2}, Ana M. Pérez García³, Jordi Cepa^{1,2}, Miguel Cerviño^{1,2,4}, and the OTELO Team




¹ Instituto de Astrofísica de Canarias, E-38205 La Laguna, Tenerife, Spain
² Departamento de Astrofísica, Universidad de La Laguna, E-38206 La Laguna, Tenerife, Spain
³ ISDEF for ESA-European Space Agency, CAB-ESAC, Camino Bajo del Castillo s/n, Villafranca del Castillo, E-28692 Villanueva de la Cañada, Spain
⁴ Instituto de Astrofísica de Andalucía - CSIC, Glorieta de la Astronomía, s/n. E-18008 Granada, Spain

OTELO in a nutshell


The OSIRIS Tunable Filter Emission Line (OTELO) survey is a 2D-spectroscopic ($R \sim 700$), blind tomography between Meinel bands in the NIR domain, on a selected 7.5×7.5 arcmin 2 field of the **Extended Groth Strip**.

The data-cube is defined by 36 slices with 12 Å of bandwidth each, sampled every 6 Å from 9280 Å towards the blue. Thus, each source accounts for a pseudo-spectrum with +36 spaxels of similar length.


Flux in each individual slice obeys the system transmission function as plotted beside. A multi-wavelength catalog with photo-z was concurrently built using ancillary data.



See the Posters of **Marina Ramón-Pérez** and **Jakub Nadolny**, or the Invited Talk of **Jordi Cepa** for more information about.



ID 6407 not only stands out as a candidate to line emitter in the corresponding C-M diagram, but it exhibits a remarkable emission feature in the OTELO's pseudo-spectrum. The line profile is consistent with predictions about the influence of the instrumental profile of tunable filters on the Ly- α line at high redshifts (de Diego et al. 2013). If the spectral feature shown corresponds to this line, **ID 6407 is a LAE at $z_{\text{spec}} = 6.531$** , which is consistent with the first photo-z solution ($z_{\text{phot}} = 6.46$). An upper limit in rest-frame EW is $\sim 18.6 \pm 5.1$ Å if continuum is actually sampled, with a moderate Ly- α luminosity of $\sim 3.49 \pm 0.9 \times 10^{42}$ erg/s.




OTELO & LAEs at $z \sim 6.5$


In the growing zoo of very high redshift sources, the **Lyman- α emitters (LAEs)** take up a singular niche. This emission is one of the most powerful tools to study galaxies during and after the reionization era.

Search of LAEs are mainly based on techniques as "dropout" (Steidel et al. 1996) or color excess in C-M diagrams, or by exploiting blind, slitless spectroscopic surveys. **OTELO is able to combine both techniques** in most cases.

Currently, we are studying **a sample of 150+ candidates** to very high-redshift galaxies in the OTELO field. A large number of interlopers is expected (i.e. cool dwarfs in the Galactic Halo, Balmer-break & other lower redshift galaxies). **An example of a LAE candidate (ID 6407) from OTELO survey is shown below.**



ID 6407 is a very faint object that subtends about 1.8 x 2.6 arcsec. Despite its low SNR, it can be seen in z (CFHTLS), OTELO and Ks (WIRDS) -bands. As expected, ID 6407 is a fair "dropout" in i-band (g+r+i image). Most LAEs are **very hard to detect in UV continuum** (Bacon et al. 2014), even in the deepest broad-band images (3.6+4.5 μm).



References:

- Bacon, R. et al. (2015) A&A 575, 75
de Diego, J.A. et al. (2013) AJ 146, 96
Steidel, C.C. et al. (1996) ApJL 462, 17